

GRAPHS THAT ADMIT A HAMILTON PATH ARE CUP-STACKABLE

PETR GREGOR, ARTURO MERINO, TORSTEN MÜTZE, AND FRANCESCO VERCIANI

ABSTRACT. Fay, Hurlbert and Tennant recently introduced a one-player game on a finite connected graph G , which they called cup stacking. Stacks of cups are placed at the vertices of G , and are transferred between vertices via stacking moves, subject to certain constraints, with the goal of stacking all cups at a single target vertex. If this is possible for every target vertex of G , then G is called *stackable*. In this paper, we prove that if G admits a Hamilton path, then G is stackable, which confirms several of the conjectures raised by Fay, Hurlbert and Tennant. Furthermore, we prove stackability for certain powers of bipartite graphs, and we construct graphs of arbitrarily large minimum degree and connectivity that do not allow stacking onto any of their vertices.

1. INTRODUCTION

In a recent paper, Fay, Hurlbert and Tennant [FHT24] introduced a one-player game on a finite connected graph G , which they called *cup stacking*. Initially, there is a single cup placed on every vertex of G . Each move consists of removing all the r cups placed on one vertex x and stacking them onto the cups of a vertex y in distance exactly r from x , provided that y already has at least one cup on it. The objective is to eventually move all cups onto a single target vertex t , in which case G is called *t -stackable*. If G is t -stackable for every vertex t , then it is called *stackable*. Note that the number of vertices with cups on them decreases by 1 with every move, so on a stackable graph the game always ends after exactly $n - 1$ moves, where n is the number of vertices of G . We remark that stackability is *not* a monotone property under adding edges. In particular, there are graphs that are stackable, which by adding edges turn into a non-stackable graph.

Cup-stacking is a recent addition to the zoo of one- and two-player games where ‘things move around’ in a finite graph, such as graph pebbling [HK21], graph pegging [HKMW09], chip firing [BLS91], cops and robbers [BN11], etc.

The paper [FHT24] proves that paths, cycles, two-dimensional grids, the d -dimensional hypercube Q_d for $d \leq 20$, Kneser graphs $K(n, k)$ for $n \geq 3k - 1$, and Johnson graphs with diameter 2 are stackable. It also characterizes which complete partite graphs are stackable. The authors also conjectured that the hypercube Q_d is stackable for all $d \geq 1$, and more generally

(Petr Gregor) DEPARTMENT OF THEORETICAL COMPUTER SCIENCE AND MATHEMATICAL LOGIC, CHARLES UNIVERSITY, PRAGUE, CZECH REPUBLIC

(Arturo Merino) UNIVERSIDAD DE O’HIGGINS, RANCAGUA, CHILE

(Torsten Mütze) INSTITUT FÜR MATHEMATIK, UNIVERSITÄT KASSEL, KASSEL, GERMANY & DEPARTMENT OF THEORETICAL COMPUTER SCIENCE AND MATHEMATICAL LOGIC, CHARLES UNIVERSITY, PRAGUE, CZECH REPUBLIC

(Francesco Verciani) INSTITUT FÜR MATHEMATIK, UNIVERSITÄT KASSEL, KASSEL, GERMANY

E-mail addresses: gregor@ktiml.mff.cuni.cz, arturo.merino@uoh.cl, tmuetze@mathematik.uni-kassel.de, francesco.verciani@uni-kassel.de.

This work was supported by Czech Science Foundation grant GA 22-15272S. The authors participated in the workshop ‘Combinatorics, Algorithms and Geometry’ in March 2024, which was funded by German Science Foundation grant 522790373.

that Cartesian products of paths of arbitrary lengths are stackable. They also asked whether all connected Kneser graphs and generalized Johnson graphs are stackable.

1.1. Our results. The main contribution of this work is to show that graphs that admit a Hamilton path are stackable (Theorem 1). This considerably enlarges the catalogue of graphs that are known to be stackable. In particular, this yields stackability of d -dimensional hypercubes for all $d \geq 1$, Cartesian products of paths of arbitrary lengths, connected Kneser graphs and generalized Johnson graphs (Corollary 2), resolving the problems raised by Fay, Hurlbert and Tennant [FHT24].

In view of this result, the remainder of our paper focusses on graphs that do not admit Hamilton paths. To this end, we present a versatile tool, which we call ‘chunking lemma’ (Lemma 5), to split paths in a bipartite graph into subpaths of the correct sizes so as to stack the cups from the subpaths to a particular target vertex. We apply this lemma to bipartite graphs with small diameter (Theorem 6), and powers of graphs (Theorem 8), including trees (Theorems 11 and 12), proving in particular that a sufficiently large power of any connected bipartite graph is stackable (Theorem 9)

Lastly, we construct graphs that are not t -stackable for any target vertex t , which we call *strongly non-stackable*. In particular, we obtain strongly non-stackable graphs with arbitrarily large minimum degree and connectivity (Theorems 15 and 16, respectively). We also construct families of graphs that are stackable, but which can be made (strongly) non-stackable by adding edges, showing that stackability is not monotone (Theorems 21 and 22). This demonstrates that our result on stackability of graphs with a Hamilton path is not a trivial consequence of the fact that paths are stackable.

1.2. Related games and rules. In our version of the game, a move of r cups from a vertex x to another vertex y requires that there is a shortest path $P(x, y)$ of length r from x to y in G . This is the reason for the non-monotonicity under adding edges, as adding an edge may change shortest paths. A similar game was studied by Veselovac [Ves22], using slightly different terminology (instead of cups being stacked, he considers frogs that jump on each other), where ‘shortest path $P(x, y)$ ’ is replaced by ‘path $P(x, y)$ ’, i.e., cups (or frogs) can move along non-geodesic paths¹, a variant of the game that was also mentioned in [FHT24] for further study. Of course, if G is stackable in the geodesic game, then G is also stackable in the non-geodesic game. Furthermore, if G is a tree both variants are identical, as any path in a tree is also a shortest path. In fact, most results in Veselovac’s thesis are about trees. Specifically, he shows that paths, stars, and so-called starfishes and dandelions are stackable. He also proves that the tree T obtained from two stars with at least 3 rays each by joining their centers with an edge is not t -stackable for any vertex t . The non-geodesic game behaves monotonically with respect to adding edges, namely if we have a stackable graph, then any graph obtained by adding edges is also stackable. In particular, since the path is stackable, any graph that admits a Hamilton path is also stackable. For the rest of this paper, we do not consider this variant of the game, but instead we focus on the game introduced by Fay, Hurlbert and Tennant [FHT24] where cups have to be moved along geodesic (i.e., shortest) paths.

Mitchell [Mit23] considers yet another variant, where ‘shortest path $P(x, y)$ ’ is replaced by ‘walk $P(x, y)$ ’, i.e., vertices and edges used during one move may repeat. As pointed out in [FHT24], other variants are of course possible, such as ‘trail $P(x, y)$ ’, i.e., vertices may repeat, but not edges.

¹The English version of the abstract of his thesis says ‘shortest paths’, but this is a mistranslation of the original Croatian version. Definition 1.2 in the thesis clearly requires ‘paths’ only, not ‘shortest paths’.

Woll [Wol17] has investigated the game on the path where in the initial configuration each vertex has either 0 or 1 cups.

1.3. Outline of this paper. In Section 2 we prove that the existence of a Hamilton path implies stackability. In Section 3 we present the chunking lemma and various applications of it to bipartite graphs and their powers (that do not admit Hamilton paths). In Section 4 we construct graphs that are not t -stackable for any of target vertex t . In Section 5 we show that stackability is not monotone under adding edges. We conclude with some open questions in Section 6.

2. GRAPHS THAT ADMIT A HAMILTON PATH

A *Hamilton path* in a graph is a path that visits every vertex exactly once.

Theorem 1. *Any graph that admits a Hamilton path is stackable.*

This substantially enlarges the catalogue of known stackable graphs by all graphs that are known to have a Hamilton path. The following corollary lists only those explicitly conjectured or mentioned in [FHT24]. To state the result, we write P_ℓ for the path with ℓ vertices. Furthermore, the *Cartesian product* $G \square H$ of two graphs $G = (V, E)$ and $H = (W, F)$ has the vertex set $V \times W$ and an edge $((v, w), (v', w'))$ whenever $v = v'$ and $(w, w') \in F$ or $(v, v') \in E$ and $w = w'$. The *d -dimensional hypercube* Q_d is the graph $\square_{i=1}^d P_2$. It is easy to show by induction that Cartesian products of paths of arbitrary lengths (see (1) below), in particular Q_d , admit a Hamilton path. For integers $k \geq 1$ and $n \geq 2k + 1$, the *Kneser graph* $K(n, k)$ has as vertices all k -element subsets of $\{1, \dots, n\}$, and edges between disjoint sets. The *generalized Johnson graph* $J(n, k, s)$ has as vertices all k -element subsets of $\{1, \dots, n\}$, and an edge between any two sets A and B that satisfy $|A \cap B| = s$. To ensure that the graph is connected, we assume that $s < k$ and $n \geq 2k - s + \mathbf{1}_{[s=0]}$, where $\mathbf{1}_{[s=0]}$ denotes the indicator function that equals 1 if $s = 0$ and 0 otherwise. Clearly, we have $J(n, k, 0) = K(n, k)$. All these graphs were recently shown to have a Hamilton cycle in [MMN23]. In fact, the Petersen graph $K(5, 2) = J(5, 2, 0) = J(5, 3, 1)$ does not have a Hamilton cycle, but a Hamilton path.

Corollary 2. *The following families of graphs are stackable:*

- *d -dimensional hypercubes Q_d for all $d \geq 1$;*
- *Cartesian product of paths $P_{\ell_1} \square P_{\ell_2} \square \dots \square P_{\ell_d}$ for all $d \geq 1$ and all integers $\ell_1, \dots, \ell_d \geq 1$;*
- *Kneser graphs $K(n, k)$ for all $k \geq 1$ and $n \geq 2k + 1$;*
- *generalized Johnson graphs $J(n, k, s)$ for all $k \geq 1$, $0 \leq s < k$, and $n \geq 2k - s + \mathbf{1}_{[s=0]}$.*

Theorem 1 is an immediate consequence of the following lemma.

Lemma 3. *Let G be a graph and $P = (x_1, \dots, x_\ell)$ a subpath of G . Then for any $t \in \{1, \dots, \ell\}$, we can stack the ℓ cups from P onto the vertex x_t .*

We write $d(x, y)$ for the distance between vertices x and y in G .

Proof. We argue by induction on ℓ . The induction basis $\ell = 1$ is trivial. For the induction step let $\ell \geq 2$. Our task is to stack the ℓ cups from $P = (x_1, \dots, x_\ell)$ onto the target vertex x_t . We assume w.l.o.g. that $t \geq 2$, otherwise we can do the argument with the reversed path. We define $s := d(x_1, x_t)$, and we split P into two paths $P' := (x_1, \dots, x_s)$ and $P'' := (x_{s+1}, \dots, x_\ell)$. Note that $s \leq t - 1$ and therefore $x_t \in P''$. By induction, we can stack the s cups from P' onto x_1 . Similarly, by induction, we can stack the $\ell - s$ cups from P'' onto x_t . In the last step, because of $d(x_1, x_t) = s$, we can move the s cups from x_1 to x_t . This completes the proof. \square

Note that this proof does not require that P is an isometric path in G , i.e., the distances between vertices on P along the path need not be the same as the distances between those vertices in the whole graph G .

By Theorem 1, a graph is stackable if it has a Hamilton path. The converse implication, however, is not true in general. The smallest graphs that are stackable but do not admit Hamilton paths have 6 vertices, and they are all shown in Figure 1.




FIGURE 1. All smallest stackable graphs that do not admit Hamilton paths.

3. BIPARTITE GRAPHS

In this section we establish stackability of various bipartite graphs, in particular graph powers, that do not have Hamilton paths; see Remark 7 below.

We will use the following important property of bipartite graphs, stated without proof.

Lemma 4. *Let G be a bipartite graph, let t be one of its vertices, and let xy be an edge of G . Then we have $d(x, t) = d(y, t) \pm 1$.*

3.1. Chunking lemma. For integers $a \leq b$ we define $[a, b] := \{a, a + 1, \dots, b\}$. We also write $[N] := [1, N] = \{1, \dots, N\}$. Given a sequence x_1, \dots, x_N of integers, a *chunk* is a consecutive subsequence x_i, x_{i+1}, \dots, x_j for $1 \leq i \leq j \leq N$, and we refer to the quantity $j - i + 1$ as its *length*. Such a chunk is *proper* if the value of one of its element equals its length, i.e., $x_k = j - i + 1$ for some $k \in [i, j]$. For example, the sequence 6, 5, 3, 4, 5, 6, 4 can be partitioned into the chunks 6, 5, 3 and 4, 5, 6, 4, both of which are proper, and this is in fact the only way to split this sequence into proper chunks.

Lemma 5. *Let x_1, \dots, x_N be a sequence of positive natural numbers such that $(x_1, x_2) \neq (2, 1)$ and $x_{i+1} = x_i \pm 1$ for $i = 1, \dots, N - 1$. If $N \geq (\max_{i=1}^N x_i)^2$, then the sequence can be partitioned into proper chunks.*

The assumption $(x_1, x_2) \neq (2, 1)$ is needed, because the sequence 2, 1, 2, 1, \dots , 2, 1, 2 cannot be partitioned into proper chunks.

We will apply Lemma 5 to partition a path P with N vertices in a bipartite graph G into subpaths. The number x_i is the distance of the i th vertex on P to some fixed target vertex t not on the path, and by Lemma 4 these numbers alternate by ± 1 by the assumption that G is bipartite. The proper chunks correspond to subpaths of P with the property that the number of vertices on the subpath equals the distance of one vertex s on the subpath to t . We can then stack all cups from the subpath onto s by Lemma 3, and from there with one additional move onto t . The crucial condition stated in the lemma for this to work is that N is at least quadratic in the largest of the distances x_i .

Proof. We say that $i \in [N]$ is a *cut* if x_1, \dots, x_i can be partitioned into proper chunks. Our goal is to prove that N is a cut.

By the assumption that $(x_1, x_2) \neq (2, 1)$ and $x_2 = x_1 \pm 1$, there are two consecutive cuts $\min\{x_1, x_2\}, \max\{x_1, x_2\}$.

Consider a sequence of $s \geq 2$ consecutive cuts $[i - s, i - 1]$. The different cases in the following proof are illustrated in Figure 2. We distinguish two main cases: If $x_i \leq s$, then we define $t := x_i$




FIGURE 2. Illustration of the proof of Lemma 5.

and we have the proper chunk $x_{i-(t-1)}, \dots, x_i$ of length t , implying that i is a cut, i.e., we have found $s + 1$ consecutive cuts $[i - s, i]$. It remains to consider the case $x_i > s$. We define $t := x_i$ and consider the proper chunks $x_{i-r}, \dots, x_{i-r+t-1}$, each of length t , for $r = 0, \dots, s - 1$, i.e., we have found s consecutive cuts $[i - s + t, i + t - 1]$. If $x_{i+1} = x_i + 1$, then we have the proper chunk x_i, \dots, x_{i+t} of length $t + 1 = x_{i+1}$, and thus another cut $i + t$. If $x_{i+1} = x_i - 1$ and $t > s + 1$, then we have the proper chunk $x_{i-s+1}, \dots, x_{i-s+t-1}$ (note that $i - s + t - 1 \geq i + 1$ by the additional assumption $t > s + 1$) of length $t - 1 = x_{i+1}$, and thus another cut $i - s + t - 1$. Otherwise we have $x_{i+1} = x_i - 1$ and $t = s + 1$. Consider the smallest $r \geq 1$ such that $x_{i+1+r} = x_{i+r} + 1$. We then have $x_i, x_{i+1}, \dots, x_{i+1+r} = s + 1, s, s - 1, \dots, s - r + 2, s - r + 1, s - r + 2$ and therefore $r \leq s$. We distinguish the subcases $r = 1$ and $r \geq 2$. If $r \geq 2$ we consider the proper chunk $x_{i+r}, \dots, x_{i+s+1}$ (note that $i + 2 \leq i + r \leq i + s$ by the additional assumption $r \geq 2$) of length $s - r + 2 = x_{i+1+r}$, which gives another cut $i + s + 1 = i + t$. It remains to consider the case $r = 1$, i.e., $x_{i+2} = x_i = x_{i+1} + 1 = s + 1$. If $x_{i+3} = x_{i+2} + 1 = s + 2$, then consider the proper chunk x_i, \dots, x_{i+s+1} of length $s + 2 = x_{i+3}$, which gives another cut $i + s + 1 = i + t$. If $x_{i+3} = x_{i+2} - 1 = s$, then consider the proper chunk $x_{i+2}, \dots, x_{i+s+1}$ of length $s = x_{i+3}$, which gives another cut $i + s + 1 = i + t$.

We define $C := \max_{i=1}^N x_i$. By the arguments from before, we see at least 2 consecutive cuts in the interval $[1, C]$. Furthermore, if there are $s \geq 2$ consecutive cuts in the interval $[1, i]$, then there are at least $s + 1$ consecutive cuts in the interval $[1, i + C + 1]$. We conclude that there are C consecutive cuts in the interval $[1, (C - 1) \cdot (C + 1)]$. Note that every number greater than this sequence of C consecutive cuts is also a cut, in particular $N \geq C^2 > (C + 1)(C - 1)$ is a cut. This completes the proof. \square

3.2. Bipartite graphs with small diameter. We refer to a spanning forest of paths in G as a *path partition* of G (the paths need not be induced). For a path P in G between vertices x and y , we define $T(P) := \{x, y\}$ as the set of terminal vertices of P . The *diameter* of G is the maximum of $d(x, y)$ over all pairs of vertices x, y in G .

Theorem 6. *Let G be a connected bipartite graph with diameter d and let t be one of its vertices. If G has a partition into $p \geq 2$ paths P_1, \dots, P_p such that $t \in P_1$ and P_j has at least d^2 vertices and $d(T(P_j)) \notin \{2, 4\}$ for all $j \in [2, p]$, then G is t -stackable.*

In this theorem and its proof, and also in Theorem 8 and its proof below, P_j , $j \in [p]$, are paths of arbitrary lengths, i.e., the index j does not refer to the number of vertices.

Proof. By Lemma 3, the cups from P_1 can be stacked onto t . Now consider one of the remaining paths P_j , $j \in [2, p]$, and let x_1, \dots, x_N be the sequence of distances from the vertices along this path to t . As $t \notin P_j$ we have $x_i \geq 1$ for all $i \in [N]$, and as G is bipartite we have $x_{i+1} = x_i \pm 1$ for all $i \in [N - 1]$ by Lemma 4. Furthermore, note that if $d(T(P_j)) \geq 5$, then we have $\max\{x_1, x_N\} \geq 3$ and we can assume w.l.o.g. that $x_1 \geq 3$, in particular $x_1 \neq 2$. On the other hand, if $d(T(P_j)) \in \{1, 3\}$, then x_1 and x_N have opposite parity by Lemma 4, in particular $x_1 \neq x_N$, so we can again assume w.l.o.g. that $x_1 \neq 2$. The remaining cases $d(T(P_j)) \in \{2, 4\}$ are excluded by the assumptions in the lemma. Lastly, as the diameter of G is d the assumption $N \geq d^2$ ensures that $N \geq (\max_{i=1}^N x_i)^2$. Therefore, the conditions of Lemma 5 are satisfied, and we obtain that the path P_j can be split into q subpaths Q_1, \dots, Q_q such that the number of vertices of Q_k equals $d(s_k, t)$ for some vertex $s_k \in Q_k$ for all $k \in [q]$. We can thus apply Lemma 3 for each path Q_k , $k \in [q]$, to stack the cups from Q_k onto s_k , and from there with one additional move onto t . \square




FIGURE 3. Illustration of how Theorem 6 applies to certain subgraphs of a biwheel.

We proceed to give an example of how to apply Theorem 6. For any integer $\ell \geq 2$ we define the *biwheel* W_ℓ as the graph with vertex set $\{c\} \cup \{x_1, \dots, x_\ell\} \cup \{y_1, \dots, y_\ell\}$ and edge set $\{(c, x_i) \mid i \in [\ell]\} \cup \{(x_i, y_i), (y_i, x_{i+1}) \mid i \in [\ell]\}$, where indices are considered modulo ℓ ; see Figure 3 (a). For any set $I \subseteq [\ell]$ we define $W_{\ell, I} := W_\ell \setminus \{(x_i, y_i) \mid i \in I\}$, i.e., the edges (x_i, y_i) for $i \in I$ are removed from the biwheel; see Figure 3 (b). We aim to show that $W_{\ell, I}$ is stackable provided that for any two integers $j, j' \in I$ we have $|j - j'| \geq 8$ (modulo ℓ). First note that the graph $W_{\ell, I}$ is bipartite and has diameter $d = 4$. Also note that $W_{\ell, I}$ has $p := |I|$ vertices of degree 1, so it does not have a Hamilton path for $p \geq 3$, i.e., Theorem 1 does not apply. Clearly, the edges of $W_{\ell, I}$ not incident with the central vertex c form a set \mathcal{P} of paths whose end vertices lie in different partition classes and hence satisfy $d(T(P)) \notin \{2, 4\}$ for all $P \in \mathcal{P}$; see Figure 3 (c). Also, each of the paths has at least $16 \geq d^2$ vertices by the earlier assumption on I . For any target vertex t of $W_{\ell, I}$, we can take P_1 to be the path from \mathcal{P} that contains t extended by the central vertex c , or if $t = c$ we extend any of the paths by c to become P_1 , and we take P_2, \dots, P_p to be the remaining paths in the set \mathcal{P} ; see Figure 3 (d). Applying Theorem 6 yields that $W_{\ell, I}$ is t -stackable, as desired.

3.3. Powers of bipartite graphs. We now apply Lemma 5 to powers of bipartite graphs G . The *r th power* of a graph G is the r -fold Cartesian product with itself, namely $G^r := \underbrace{G \square G \square \dots \square G}_{r \text{ times}}$.

Remark 7. For a bipartite graph G , we write $\Delta(G) \geq 0$ for the difference in size between the partition classes of G . Note that G^r is bipartite, and furthermore satisfies $\Delta(G^r) = \Delta(G)^r$. Consequently, if $\Delta(G) \geq 2$, then $\Delta(G^r) \geq 2^r \geq 2$, and so G^r does not admit a Hamilton path for any $r \geq 1$. The results below thus yield infinitely many graphs that are stackable, but that do not admit a Hamilton path (in a very strong sense the larger $\Delta(G^r)$ is; so Theorem 1 does not apply).

Theorem 8. *Let G be a connected bipartite graph with diameter d . If G has a partition into $p \geq 2$ paths P_1, \dots, P_p with at least $k \geq 2$ vertices each, then for any integer $r \geq 2$ that satisfies $k^r \geq (dr)^2$ the graph G^r is stackable.*

The Cartesian product of paths $P_1 \square \dots \square P_p$, where $P_p = (x_1, x_2, \dots)$, admits a *canonical Hamilton path* $H(P_1, \dots, P_p)$, defined recursively as $H(P_1) := P_1$ if $p = 1$ and

$$H(P_1, \dots, P_p) := H'x_1, \text{rev}(H')x_2, H'x_3, \text{rev}(H')x_4, \dots, \quad \text{if } p \geq 2, \quad (1)$$

where $H' := H(P_1, \dots, P_{p-1})$ and $\text{rev}(H')$ is the path H' traversed in reverse order. For example, for $P_1 = (a, b, c)$ and $P_2 = (d, e, f)$ we have

$$H(P_1, P_2) = (a, d), (b, d), (c, d), (c, e), (b, e), (a, e), (a, f), (b, f), (c, f).$$

Proof of Theorem 8. For $i \in [p]$ we write $N_i \geq k \geq 2$ for the number of vertices of the path P_i . The graph G^r is bipartite, has diameter dr , and it has an induced partition into r -dimensional grids $Q(i_1, \dots, i_r) := P_{i_1} \square P_{i_2} \square \dots \square P_{i_r}$ for integers $i_1, \dots, i_r \in [p]$. Each of the grids $Q(i_1, \dots, i_r)$ admits a canonical Hamilton path and contains $\prod_{j=1}^r N_{i_j}$ vertices. Fix an arbitrary vertex $t = (t_1, \dots, t_r)$ of G^r . By Lemma 3, we can stack the cups from $Q = Q(i_1, \dots, i_r)$ with $t \in Q$ onto t . Now consider one of the remaining grids $Q = Q(i_1, \dots, i_r)$ with $t \notin Q$. We assume w.l.o.g. that $t_1 \notin P_{i_1}$. Furthermore, for each $j \in [2, r]$ we choose the orientation of the path P_{i_j} so that t_j is not its first vertex (this is possible because it has $N_{i_j} \geq 2$ vertices), and we let $x_{j,1}, \dots, x_{j,N_{i_j}}$ be the sequence of distances from the vertices along this path to t_j in the graph G . By the assumption that $t_1 \notin P_{i_1}$ we have $x_{1,k} \geq 1$ for all $k \in [N_{i_1}]$, and by the choice of orientation of P_{i_j} we have $x_{j,1} \geq 1$ for all $j \in [2, r]$. Let x_1, \dots, x_N , $N := \prod_{j=1}^r N_{i_j} \geq k^r$, be the sequence of distances from the vertices along the canonical path $H = H(P_{i_1}, \dots, P_{i_r})$ to t in the graph G^r . As $t \notin Q$ we have $x_i \geq 1$ for all $i \in [N]$, and as G^r is bipartite we have $x_{i+1} = x_i \pm 1$ for all $i \in [N-1]$ by Lemma 4. Furthermore, note that $x_1 = \sum_{j=1}^r x_{j,1}$, which if $r \geq 3$ implies that $x_1 \geq 3$ and in particular $x_1 \neq 2$. If $r = 2$ and $x_1 = 2$, then we have $x_{1,1} = x_{2,1} = 1$ and therefore $x_{1,2} = 2$, implying that $x_2 = x_{1,2} + x_{2,1} = 3$ and therefore $(x_1, x_2) \neq (2, 1)$. Lastly, as the diameter of G^r is dr the assumption $N \geq k^r \geq (dr)^2$ implies that $N \geq (\max_{i=1}^N x_i)^2$. Therefore, the conditions of Lemma 5 are satisfied, we can split the path H into subpaths, stack the vertices from each subpath onto one of its vertices that has the correct distance to t by Lemma 3, and from there with one additional move onto t . \square

Corollary 9. *For any connected bipartite graph G that has a partition into $p \geq 2$ paths with at least $k \geq 2$ vertices, there is an integer $r_0 \geq 2$ such that G^r is stackable for all $r \geq r_0$.*

Proof. Let d be the diameter of G . Note that k^r grows exponentially with r , whereas $(dr)^2$ grows quadratically with r , so there is an integer r_0 such that $k^r \geq (dr)^2$ for all $r \geq r_0$, and then Theorem 8 applies. \square

3.4. Powers of trees. To find a partition of G into long paths, it suffices to find a spanning tree of G in which any two leaves have large distance. Note that the diameter of a tree T is the maximum of $d(x, y)$, taken over all pairs of leaves x and y . We define the *spread* of a tree T as the minimum of $d(x, y)$, taken over all pairs of distinct leaves x and y .

Lemma 10. *Let T be a tree with spread k . Then T has a partition into paths on at least $k/2$ vertices each.*

Proof. This proof is illustrated in Figure 4. For any two vertices x and y in T , we write $P(x, y)$ for the path from x to y in T . Consider T rooted at one of its leaves r . The partition of T into paths is obtained by repeatedly removing a path from this rooted tree as follows: We choose a leaf x at maximum distance from r , let y be the vertex on $P(r, x)$ farthest from r that has more than one descendant, and we let y' be the descendant of y on $P(r, x)$. The path $P(y, x)$ has length at least $k/2$, otherwise x would have distance less than k from one of the other descendant leaves of y . It follows that $P(y', x)$ has at least $k/2$ vertices. So we remove $P(y', x)$ from T as part of the partition and repeat the process, with the same root r , until there is only one path left (this last path has length at least k and therefore at least $k+1$ vertices). As removing $P(y', x)$ from the tree does not create additional leaves, this yields the desired partition. \square




FIGURE 4. Illustration of the proof of Lemma 10. Part (a) shows one step of the partition, and (b) shows the final partition of T into paths, where the numbers $1, \dots, 9$ indicate the order in which the paths are removed.

Theorem 11. *Let T be a tree with spread $k \geq 72$ and diameter at most $k^{3/2}/\sqrt{72}$. Then T^3 is stackable.*

Proof. As T is a tree, it is clearly bipartite. Furthermore, by Lemma 10, T admits a partition into paths with at least $k/2 \geq 36$ vertices each. For $d := k^{3/2}/\sqrt{72}$ and $r := 3$ the inequality $(k/2)^r \geq (dr)^2$ is satisfied, as shown by the following computation:

$$(k/2)^3 \geq ((k^{3/2}/\sqrt{72}) \cdot 3)^2 \iff 1/8 \geq 9/72 = 1/8.$$

The claim hence follows from Theorem 8. \square

The above proof works in fact also for $k \geq 3$, but the statement of the theorem would be void for $k = 3, \dots, 71$, because in those cases $k > k^{3/2}/\sqrt{72}$, and the diameter of any tree is always greater or equal to the spread.

The *s-subdivision* of a graph G is the graph obtained by replacing every edge of G by a path of length s .

Theorem 12. *Let T be a tree with spread k and diameter d . Let T' be the s -subdivision of T for some $s \geq 72d^2/k^3$. Then $(T')^3$ is stackable.*

Proof. We want to apply Theorem 11 to T' . Note that T' has spread $k' = sk$ and diameter $d' = sd$, so the inequality $d' \leq k'^{3/2}/\sqrt{72}$ is equivalent to $sd \leq (sk)^{3/2}/\sqrt{72}$, which is equivalent to the assumption $s \geq 72d^2/k^3$. Note also that $k' = sk \geq 72$ by the assumption $s \geq 72d^2/k^3$ and the observation that $d \geq k$. \square

4. STRONGLY NON-STACKABLE GRAPHS

In this section we construct families of graphs G that are not t -stackable for *any* target vertex t of G . Recall that we refer to such a graph as *strongly non-stackable*. Recall that Veselovac [Ves22] proved that the tree obtained from two stars with at least 3 rays by joining their centers with an edge is strongly non-stackable. In this section, we enlarge the catalogue

of strongly non-stackable graphs. In particular, we show that there are strongly non-stackable graphs with arbitrarily large minimum degree and connectivity (which must be non-trees).

The next result is a versatile result to prove non-stackability based on the presence of large independent sets in the graph.

Lemma 13. *Let G be a connected graph with vertex set V and let $t \in V$. Let U be an independent set in G , and define $U' := \{x \in U \mid d(x, t) \geq 2\}$, $W := V \setminus U$ and $d := \max_{x \in V} d(x, t)$. If $|U'| > (d - 1)|W|$, then G is not t -stackable.*

In words, the set U' are those vertices from the independent set U that have distance at least 2 from t , the set W is the complement of U , and d is the maximum distance of vertices from t . The inequality $|U'| > (d - 1)|W|$ expresses that there are many more vertices in U' than in W . Intuitively, the graph G referred to in Lemma 13 has small diameter, and a large independent set, a large subset of which are non-neighbors of t .

Proof. For the sake of contradiction suppose that G is t -stackable. Consider one of the cups placed on a vertex $x \in U'$. As $d(x, t) \geq 2$, this cup cannot be moved to t directly, but it is moved to t in a sequence of moves, where the final move involves $c \geq 2$ cups being moved from some vertex to t . Consider the origin of these c cups. One of them originated from x by construction. Furthermore, not all of these c cups can originate from vertices in U , simply because any two vertices in U have distance at least 2, so the first move in the sequence had to involve a vertex from W , which is empty after this sequence of moves. It follows that this sequence of moves emptied at least one vertex from W and at most $c - 1$ vertices from U' . As $|U'| > (d - 1)|W|$ and $d \geq c$ by the definition of d , we conclude that at the end of the game, at least one of the vertices in U' still has a cup on it, a contradiction. \square

Given a graph G , the *c -cactus of G* is the graph obtained from G by attaching c pendant edges to each vertex of G ; see Figure 5.

Theorem 14. *Let G be a connected graph with $n \geq 2$ vertices and diameter d , and consider the c -cactus G' of G for $c > (d + 1)n/(n - 1)$. Then G' is strongly non-stackable.*

Figure 5 shows examples of applying Theorem 14 to construct strongly non-stackable graphs.




FIGURE 5. Three examples of the cactus construction in Theorem 14. The graphs G' on the right hand side are strongly non-stackable, i.e., not t -stackable for any vertex t .

Proof. Let V be the vertex set of G . We can take all degree 1 vertices of G' as the set U , i.e., $|U| = cn$ and $|W| = |V \setminus U| = n$. For any vertex t of G' we have $|U'| \geq c(n - 1)$. Furthermore, the diameter of G' equals $d' := d + 2$. Consequently, the inequality $|U'| > (d' - 1)|W|$ is satisfied, as $c(n - 1) > (d + 1)n$ is equivalent to the assumption $c > (d + 1)n/(n - 1)$. \square

Generalizing this construction, given any graph G , we can glue copies of a large enough complete bipartite graph $K_{c,c}$ to each vertex of G , which produces a strongly non-stackable graph with arbitrarily large minimum degree. This yields the following result.

Theorem 15. *For any integer $c \geq 1$, there is a strongly non-stackable graph G with minimum degree c .*

This can be strengthened as follows.

Theorem 16. *For any integer $c \geq 1$, there is a strongly non-stackable graph G that is c -connected.*

Proof. Such a graph G can be constructed as follows: Take a copy of $K_{c,c}$, and to each of its partition classes, glue one copy of $K_{c,5c}$ (with its smaller partition class). We can take all degree c vertices of G' as the set U , i.e., $|U| = 10c$ and $|W| = 2c$. For any vertex t of G we have $|U'| \geq 5c$. Furthermore, the diameter of G' equals $d' = 3$. Consequently, the inequality $|U'| > (d' - 1)|W|$ is satisfied, as $5c > 2 \cdot 2c$ is equivalent to $5 > 4$. \square




FIGURE 6. Construction of c -connected strongly non-stackable graphs explained in the proof of Theorem 16. Part (a) shows the general construction, and (b) two concrete examples for $c = 1$ and $c = 2$.

In view of Lemma 13, one may wonder whether there are strongly non-stackable graphs with small independent sets. Theorem 18 below shows that the answer is yes, and the construction is based on the following lemma.

Lemma 17. *Let G be a graph with diameter at most 3 that has two degree-1 vertices u, v with a common neighbor, and let t be a target vertex in distance 2 from u and v . Then G is not t -stackable.*

Proof. Let w be the common neighbor of u and v . The cups on u and v cannot move to t directly, as they have distance 2 from t . Consequently, each of them moves to t in a stack of size 2 or 3 (as the diameter is at most 3). A stack of size 2 formed with a cup from u or v has to use the cup on w . Similarly, a stack of size 3 formed with both cups from u and v has to use the cup on w . However, a stack of size 3 located on u, v or w cannot move to t , as u, v, w have distance 2 or 1 from t . It can also not move anywhere else, as this would make it a stack of size at least 4,

and then it could not move anymore. Consequently, not both cups on u and v move to t in the same stack, and at least one of them, w.l.o.g., u say, moves in a stack that does not contain the cup from w . However, this stack must again have size 3 and must be located on u (as the single cup from u cannot move across w), which again means that this stack can never move to t . \square

The family of graphs referred to in the next theorem is illustrated in Figure 7.

Theorem 18. *Let G be a graph obtained from a clique of size at least 2 by attaching groups of at least three pendant edges to at least two of the clique vertices. Then G is strongly non-stackable.*

Proof. G has diameter 3, and regardless of the choice of target vertex t in G , there are always two degree 1 vertices with a common neighbor in distance 2 from t , so Lemma 17 applies. \square




FIGURE 7. Illustration of the graphs in Theorem 18.

5. STACKABILITY IS NOT MONOTONE

In this section, we show that the property of being stackable is not monotone under adding edges to the graph.

Fay, Hurlbert and Tennant [FHT24] completely characterized stackability of complete partite graphs. We need the following special case of their more general result. In this statement, $K_{a,b}$ is the complete bipartite graph with partition classes of size a and b .

Theorem 19 ([FHT24, Cor. 12]). *The graph $K_{a,b}$, $a \leq b$, is stackable if and only if $b = a$ or $b = a + 1$. The graph $K_{a,b}$, $a + 2 \leq b$ is not t -stackable for any vertex t in the larger partition class, but it is t -stackable for any vertex t in the smaller partition class.*

We note that the non-stackability statement in the previous theorem can be proved by applying Lemma 13 with U as the larger partition class.

Veselovac [Ves22] proved the following result. Let F_n be the graph formed from a path on $n - 2$ vertices by appending two pendant edges to the same end vertex of the path; see Figure 8.

Theorem 20 ([Ves22, Thm 3.10]). *The graph F_n is stackable for all $n \geq 9$.*

Theorem 21. *There are infinitely many graphs $G \subseteq H$ on the same vertex set such that G is stackable but H is non-stackable.*

Proof. We take $G := F_n$ for some even integer $n \geq 10$, which is stackable by Theorem 20. The graph G is bipartite with partition classes of size $n/2 - 1$ and $n/2 + 1$. Consequently, it is a subgraph of $K_{n/2-1, n/2+1}$, which by Theorem 19 is non-stackable. \square

The previous proof argues that F_{10} is stackable, whereas $K_{4,6} \supset F_{10}$ is not. In fact, Figure 8 shows a sequence of five edges e_1, \dots, e_5 of $K_{4,6} \setminus F_{10}$ that can be added to F_{10} one after the other such that the resulting graphs are alternately stackable and non-stackable, respectively, which has been checked by computer.




FIGURE 8. Illustration of the non-monotonicity of stackability under adding edges. For the non-stackable graphs, the vertices t for which the graph is not t -stackable are highlighted.

Theorem 22. *There are two graphs $G \subseteq H$ on the same vertex set such that G is stackable but H is strongly non-stackable.*

Proof. Such a pair of graphs G and H on 17 vertices is shown in Figure 9. Specifically, H is obtained from a clique on 11 vertices by appending two triples of pendant edges to two of its vertices, and G is obtained by removing edges from the clique so that it becomes a path between the appendix vertices. The graph G is stackable as shown in Figure 10, whereas H is strongly non-stackable by Theorem 18. \square




FIGURE 9. Illustration of the graphs in Theorem 22.

6. OPEN QUESTIONS

We conclude with some challenging open questions.

Can we find other interesting applications for our chunking lemma (Lemma 5)?

Can stackable trees be characterized?

Theorems 21 and 22 show that stackability is not monotone under adding edges. How ‘bad’ can this become? Specifically, can we find, for each integer $r \geq 3$, a sequence of graphs $G_1 \subseteq G_2 \subseteq \dots \subseteq G_r$ on the same vertex set, such that G_1, G_3, G_5, \dots are stackable and




FIGURE 10. Stackability of the graph G from Theorem 22. By symmetry, only the 7 target vertices marked by the little squares need to be checked. Solutions on subpaths or -stars (highlighted by colors) are not shown in full detail for clarity.

G_2, G_4, G_6, \dots are (strongly) non-stackable? Furthermore, are there graphs $G \subseteq H$ on the same vertex set that differ only in few edges (ideally only one), such that G is stackable and H is strongly non-stackable?

We may also consider a refined version of the game, where the *weight* of a move is the number of cups that are being moved. For a given graph G and target vertex t , we are looking for the sequence of moves with target vertex t that minimizes the sum of weights of all moves, and we write $\mu(G, t)$ for this quantity. We can think of $\mu(G, t)$ as the total weight of the cups being ‘transported’ during a solution. Table 1 shows this quantity for all target vertices of paths P_n on vertices $1, \dots, n$ for all $n \leq 12$. What can we say about $\mu(G, t)$ for interesting stackable graphs G ?

REFERENCES

- [BLS91] A. Björner, L. Lovász, and P. W. Shor. Chip-firing games on graphs. *European J. Combin.*, 12(4):283–291, 1991.
- [BN11] A. Bonato and R. J. Nowakowski. *The game of cops and robbers on graphs*, volume 61 of *Student Mathematical Library*. American Mathematical Society, Providence, RI, 2011.
- [FHT24] P. Fay, G. Hurlbert, and M. Tennant. Cup stacking in graphs. *Discrete Math.*, 347(9):Paper No. 114097, 13 pp., 2024.
- [HK21] G. Hurlbert and F. Kenter. Graph pebbling: a blend of graph theory, number theory, and optimization. *Notices Amer. Math. Soc.*, 68(11):1900–1913, 2021.
- [HKMW09] G. Helleloid, M. Khalid, D. P. Moulton, and P. M. Wood. Graph pegging numbers. *Discrete Math.*, 309(8):1971–1985, 2009.
- [Mit23] J. Mitchell. Jumping frogs on cyclic graphs. Master’s thesis, Murray State University, 2023.

TABLE 1. Minimum weight solutions on paths with $n = 1, \dots, 12$ vertices.

n	$\mu(P_n, i), i = 1, \dots, n$
1	{0}
2	{1, 1}
3	{3, 2, 3}
4	{4, 4, 4, 4}
5	{6, 5, 6, 5, 6}
6	{9, 7, 7, 7, 7, 9}
7	{11, 10, 9, 8, 9, 10, 11}
8	{12, 12, 12, 10, 10, 12, 12, 12}
9	{14, 13, 14, 13, 12, 13, 14, 13, 14}
10	{17, 15, 15, 15, 15, 15, 15, 15, 15, 17}
11	{19, 18, 17, 16, 17, 18, 17, 16, 17, 18, 19}
12	{22, 20, 20, 18, 18, 20, 20, 18, 18, 20, 20, 22}

[MMN23] A. Merino, T. Mütze, and Namrata. Kneser graphs are Hamiltonian. In *STOC'23—Proceedings of the 55th Annual ACM Symposium on Theory of Computing*, pages 963–970. ACM, New York, [2023] ©2023.

[Ves22] M. Veselovac. Frog jumping problem on simple graphs. Master’s thesis, J.J. Strossmayer University of Osijek, Croatia, 2022.

[Wol17] C. Woll. Jumping frogs—Challenge #2. https://mathpickle.com/wp-content/uploads/2017/03/Jumping_Frogs___Challenge__2.pdf, 2017.